

تختلف جرعات مسببات الحساسية بين الدراسات والبروتوكولات. أعلى جرعة يمكن تحملها هي كمية مسبب الحساسية التي يمكن تحملها دون حدوث رد عكسي سلبى<sup>1</sup>. قد يكون مثال على الجرعة القصوى التي يمكن تحملها هو 600 ملغ من مكافئ بروتين مسبب الحساسية<sup>1</sup>.



قائمة المراجع:

1. Casale TB, Haselkorn T, Ciaccio CE, Sriaroon P, Chipps BE. Harmonization of terminology for tolerated and reactive dose in food allergy immunotherapy. The Journal of Allergy and Clinical Immunology. In Practice. 2019;7(2):389-392.
2. Groetch, M. et al. (2022) "Retail food equivalents for post-oral immunotherapy dosing in the omalizumab as monotherapy and as adjunct therapy to multi-allergen oral immunotherapy in food-allergic children and adults (outmatch) clinical trial." The Journal of Allergy and Clinical Immunology: In Practice [Preprint].
3. Zuberbier, T. et al. (2021) "Proposal of 0.5mg of protein/100g of processed food as threshold for voluntary declaration of food allergen traces in processed food—a first step in an initiative to better inform patients and avoid fatal allergic reactions: A ga'len position paper," Allergy, 77(6), pp. 1736–1750.

**Genentech**

A Member of the Roche Group